



Diversity of arbuscular mycorrhizal fungi in the rhizosphere of date palm tree (*Phoenix dactylifera*) in Tafilalet and Zagora regions (Morocco)

Fadoua Sghir¹, Jihane Touati¹, Mohamed Chliyah¹, Amina Ouazzani Touhami¹, Abdelkarim Filali-Maltouf², Cherkaoui El Modafar³, Abdelmajid Moukhli⁴, Ahmed Oukabli⁵, Rachid Benkirane¹, and Allal Douira^{1*}

¹ Laboratoire de Botanique et de Protection des Plantes, UFR de Mycologie, Département de Biologie, Faculté des Sciences BP. 133, Université Ibn Tofail, Kénitra, Maroc

² Laboratoire de Microbiologie et Biologie Moléculaire, Faculté des Sciences, Université Mohammed V Agdal, Av Ibn Batouta, BP 1014 Rabat, Maroc

³ Laboratoire de Biotechnologie, Valorisation et Protection des Agroressources, Faculté des Sciences et Techniques Guéliz, B.P. 618, 40 000 Marrakech, Maroc

⁴ UR, Amélioration génétique des plantes, Institut national de la Recherche agronomique F- 40 000 Marrakech, Maroc

⁵ Institut national de la Recherche agronomique, Amélioration des plantes et Conservation des ressources phytogénétiques CRRRA, BP 578, Meknès, Maroc

E-mail: douiraallal@hotmail.com

Abstract

A study of arbuscular mycorrhizal fungi in the rhizosphere of date palm was conducted in the palm groves of Tafilalet and Zagora (southeastern Morocco). The parameters considered are the root colonization of date palm, spore density and species richness. The average frequencies and intensities of colonization are 66% and 7.34% respectively. The spore density varies between 80% and 132 spores / 100 g of soil, according to the site. Nine species of arbuscular mycorrhizal fungi were identified in all study sites; their appearance frequency is from 1 to 6%. Species richness varies from 2 to 7 species depending on the sites.

Keywords: Morocco, *Phoenix dactylifera*, rhizosphere, arbuscular mycorrhizal fungi (AMF), diversity, mycorrhizal parameters.

INTRODUCTION

The date palm (*Phoenix dactylifera* L.), perennial monocotyledon, is an essential species in the oasis ecosystem (Zougari-Elwedi et al., 2012). It protects the oasis against desert influences and creates a microclimate for the installation of other cultures underlying (Ben Hamida, 2011). In Morocco, the date palm is very important from an economic and human term, it contributes from 40 to 60% of revenues for one million inhabitants (Haddouch, 1993, Lambert, 2003). Indeed, the date palm production offers the backbone of the economy of oasis farms through its significant contribution in the vegetable crude product (63%) in crop gross margin as well as the sustainability of life it offers (Sbiai, 2011).

Unfortunately, the Moroccan palm grove is threatened due to several problems affecting its effective and its diversity, the most important are the scarcity of water resources accompanying climate change, soil salinity and Fusarium wilt (the Bayoud), *Fusarium oxysporum* f. sp. *albedinis* (Zeddouk, 2005; Saaidi, 1992 ; Bouammar, 2010; Radi et al., 2011; Abohatem, 2011).

Bayoud disease or vascular wilt of date palm destroyed more than two thirds of phoenicicole heritage of the area; even more the disease has a predilection for Noble varieties (Majhool) or high value (Boufeggous) (Zeddouk, 2005). The reconstruction of Moroccan palm requires to make available to farmers a lot of vigorous and protected date palm seedlings ready for planting so they remain in an environment hostile (Senoussi, 1999)

by the constraints including bayoud disease (Killian et Maire, 1930; Jaiti, 2007). This disease, which is widespread in Morocco, constitutes also a real scourge in a large part of Algeria palm groves (Djerbi, 1982 Sedra, 2005). The threat remains inevitable since Bayoud grows and spreads easily from one area to another (Senoussi, 1999).

The application of new biotechnologies, such as the mycorrhization of date palm seedling in nurseries, can help solve the problems of biotic and abiotic stresses. Plant roots mycorrhization increases significantly the absorption of phosphorus and microelements (Strullu and Plenchette, 1991 Harrison, 1999; Labidi et al, 1970, 2011.). It improves the growth of date palm seedlings (Oihabi 1991; Radi et al, 2011.), especially in soil lacking in nutrients (Al-Waibi and Khaliel, 1994) and improves the resistance to pathogens (Bartschi et al., 1981; Oihabi, 1991; Harrier et Watson, 2004, Souna et al., 2010; Abohatem, 2011). Mycorrhization also strengthens resistance to saline and water stress (Tinker, 1975; Duddridge et al., 1980; Berta, 2005; Nehila, 2012). The use of mycorrhizal fungi requires knowledge of the diversity of these fungi in the palm groves. Worldwide, communities of fungi associated with the rhizosphere of date palm are not well known. A study in Morocco on mycorrhizal fungi has noted 10 morphotypes associated with the rhizosphere of date palm in Tafilalet oases (Bouamri et al., 1986). Another study, conducted at Soltanat Oman, revealed new taxa related to date palm (Al-Yahya'ei, 2008).

This work is a continuation of previous studies. The objective is to know better the community of mycorrhizal fungi associated with the rhizosphere of date palm grown in Moroccan oasis of Tafilalet and Zagora.

MATERIAL AND METHODS

Soil and roots sampling

Two areas were selected for soil sampling at the rhizosphere of the date palm: Errachidia (Meski and Zouala) and Zagora. In each plot (three plots / site), soil samples (about 500 g each) were collected from five date palm trees at random. The sample was taken around the trunk to a depth of 0-20 cm. Finest

roots were harvested at the same time as the ground. For each plot, five samples were mixed, forming composite samples.

Physico-chemical soil analysis

The main physico-chemical characteristics of the soil were determined by conventional analyzes performed by the laboratory analyzes of soils ORMVAG of Kenitra.

Mycorrhizal degree estimation

Root preparation

The technique used is that of Philips & Hayman (1970). The roots are washed with water and finer fragments are cut into approximately 1 cm and placed into vials containing a solution of 10% potassium hydroxide. These bottles are then placed in a water bath at 90 ° C for 15 min. The roots fragments are then bleached by adding a few drops of H₂O₂ (100V) KOH mixture for 15 min. After rinsing with distilled water, they are stained with cresyl blue (0.05%) for 15 min.

Mycorrhizal Assessment

The mycorrhiza estimation is made as described by Trouvelot et al. (1986). 30 fragments are mounted between blades and blades in glycerol at 10 fragments per blade. Each fragment was checked carefully over its entire length, at the magnification × 100 and × 400.

The proposed rating system is based on the overall assessment of each of these 30 fragments. The parameters evaluated are:

- F: frequency of infection (% of number of endomycorrhizal root fragments);
- M: intensity of infection developed in endomycorrhizal part of the root system (cortex colonized proportion, expressed in %);
- A: Arbuscular content of infection reduced to the whole root system (the proportion of the root cortex containing arbuscules, expressed in %).

Spores extraction:

Spores were extracted by the method of Walker (1982). An amount of 100 g of soil was poured into a beaker filled with water. The mixture is stirred vigorously. After 10 to 20 seconds of rest, the

supernatant is transferred to another beaker which is stirred and left to stand for 10 to 30 seconds again. The suspension is then passed through four sieves (500, 200, 80 and 50 microns) superimposed. The refusal of the sieves 200, 80 and 50 microns is collected in a 100 ml beaker. This content is moved and shared in two tubes then centrifuged for 4 min at 9000 r / min. The supernatant is discarded and the tubes are filled with sucrose and centrifuged again for 15 to 30 seconds. The supernatant is collected on a sieve of 50 microns using a water jet.

Spores number estimation

The estimate was made by counting under binocular microscope the number of spores in one ml of the supernatant and by extrapolation to the total volume (100 ml). If no spores are observed, all the supernatant was reduced to 1 ml and observed again. An attempt to identify the genus of spores was performed based on the criteria proposed by Schenck & Smith (1982).

Species richness and spores appearance frequency:




Fig. 1: Different mycorrhizal structures in fragments of date palm roots; eh: external hyphae; S: spore; SP, sporocarp; a: arbuscular; v: vesicle; fe: endophyte form of sclerotia in root cortical cell.




Fig. 2: Average mycorrhizal frequency of date palm roots in the sites studied.




Fig. 3: Average mycorrhizal intensity of date palm roots in the sites studied.




Fig. 4: Average spore density of AM fungi in the rhizosphere of date palm in the sites studied.




Fig. 5: Average vesicular content of date palm roots in the sites studied.




Fig. 6: Average arbuscular content of date palm roots in the sites studied.




Fig.7 : Fréquence d'apparition d'espèces mycorhiziennes au niveau de chaque site.




Fig. 8 : Richesse spécifique des espèces mycorhiziennes dans la rhizosphère du palmier dattier dans les sites étudiés. .



Fig.9 : Spores of :*Entrophospora spl* (A), *Entrophospora kentinensis* (B), *Acaulospora colossica* (C), *Glomus spl* (D), *Glomus clarum* (E), *Acaulospora denticulata* (F), *Entrophospora sp2* (G.), *Glomus proliferum* (H) et *Glomus sp2* (I) .

Species richness is the total number of observed species by sampling site and the species occurrence frequency is the percentage of sites where each species is detected.

Statistical Analysis

The statistical treatment of results focused on the analysis of variance to a single classification criterion (ANOVA1).

RESULTS

The physico-chemical data conducted in the laboratory of Soil Analysis of the Office of Agricultural Development Gharb (ORMVAG) show that the soil rhizosphere of date palm of surveyed regions are characterized by alkaline pH (range from 8,2, and 2 site Meski 8.29, site Zouala 3) and a relatively high conductivity particularly in Zouala site 1 (2.78 mmhos / cm). The carbon rates fluctuated between 0.63% (site Meski 2) and 1.80% (Zouala 2) and those of the inorganic nitrogen varied between 73.28 ppm (Zagoura 2) and 268.16 ppm (2 Zouala). The organic matter content does not exceed 3.11% (Zouala 2) those of assimilable phosphorus ranged from 8 ppm (Meski 2) and 61 ppm (Zouala 1). Levels of exchangeable potassium are also around 294 ppm (Meski 2 Zagoura1) and 1998 ppm (Zouala 3).

The structures of AM fungi were demonstrated in all date palm root samples collected (Fig 1.): Intracellular hyphae without partition, vesicles and / or arbuscules which indicate that it is Arum arbuscular combination type. The hyphae are thin, parallel and other with tortuous and thicker wall. Occasionally there is endophytic with septate hyphae hyaline or melanized. The vesicles are also of various shapes: oval, oblong.

The mycorrhizal frequency of date palm roots varies from one site to another (Fig. 2). It ranges from 30% (Zouala 2) and 66% (Zouala1, Meski Zagoura 1 and 3). The roots mycorrhizal intensity (Fig. 3) did not exceed 7.34% (Zouala 2). The lower intensity was observed at Zouala 1 (1.66%).

The highest vesicular content (Fig. 5) is recorded in Meski 2 (4%), while at the Zouala 3 site the vesicular content is zero. Arbuscular content (Fig.

6) varies from 1.5% in Zouala1 to 7.5 found in Meski 3.

The specific richness recorded in Zouala1, Meski2 and Meski1 is about 7 species (Fig 7.), that recorded in Zagoura 1 Zagoura 2 and Zagoura 3 is in the range of 4 species; while specific richness in Meski 3 Zouala 2 and Zouala 3 is about two species. The study of the species appearance frequency showed that *Glomus clarum* (60%) is the most dominant followed by *Glomus sp1* (20%); *Acaulospora denticulata* and *Entrophospora kentiniensis* have the lowest frequency of occurrence (1%) at all sites studied (Fig. 8).

DISCUSSION AND CONCLUSION

The presence of different mycorrhizal structures in the rhizospheric soil collected from the studied sites confirmed the mycorrhizal status of date palm, considered as a mycothrophic species (Bouamri et al., 2006). The roots of date palm were receptive to arbuscular mycorrhizal fungi (Khaliel and Abu Heilah, 1985 Oihabi 1991, Al-whaibi and Khaliel, 1994) and ectomycorrhizal fungi (Farah et al., 2012). Endophytes were also observed in the roots of the date palm: septate hyphae and sclerotia in cells of the cortical parenchyma (Junppomen, 2001, Peterson et al, 2004 and 2008.).

The medium mycorrhizal intensity and frequency obtained seem low compared to those recorded by Bouamri in Morocco (2006), On *Phoenix dactylifera* (F varies between 72 and 100% and M between 5 million and 43%).

Marschner & Cakmak (1986) reported that the presence of certain chemicals in high concentrations often induces a decrease in the mycorrhizal rate. This phenomenon has been demonstrated by Amijee et al. (1989) for phosphorus. Indeed, roots colonization by mycorrhiza is maximal when the P concentration is low, and it decreases as the concentration increases. Very low P concentrations may decrease the mycorrhizal rate (Lagrange, 2009). In Zouala site, the P content is highest (61 ppm) but the mycorrhizal intensity is very low, around 1.66%. In contrast, at the site of Meski 3, the P content is 27 ppm and the mycorrhizal intensity is 5.7%. Similarly, it was noted that for low nitrogen

content (8 ppm), site Meski 2 has a low mycorrhizal intensity (3.5%).

This negative correlation observed between the root colonization rate and phosphorus reported in various studies related to date palm (Bouamri, 2006) confirms the adaptation of AMF to low soil levels of phosphorus (Smith and Read, 1997; Kaushal, 2000; Mohammad et al, 2003.).

Variation in vesicular (storage organs) and arbuscular contents (place of contact and exchange of elements between the soil and the plant) (Pawlowska and Taylor, 2004; Rosendahl, 2008) at the studied sites may depend on the physicochemical properties of soil. The variation of soil pH, temperature and effluent pollution are among the decisive factors in the distribution of mycorrhizal fungi (Mahesh and Selvaraj, 2008).

Analysis of AMF spores communities found in the rhizosphere of date palm showed that on average their number does not exceed 132 spores / 100 g of soil. The abundance of spores recorded is very low compared to that found by Bouamri (2006) in Tafilalet's soils (2080 spores/100g of soil). Work conducted on the date palm rhizosphere of Saudi Arabia (Mohamed et al., 2010) has shown that the density of spores was 58.3 to 82.3 spores /15 g of soil (from 388.66 to 548 66/100 g of soil).

This low density of spores may be due to the microclimate (Koske, 1987), the physicochemical and microbiological properties of the soil (Anderson et al, 1984. Johnson et al, 1991.) And also to the sampling season (Gemma et al. 1989). Sieverding (1991) reported the influence of nutrient levels on the spore's density.

In general, the results showed that there is no relationship between the number of spores and root intensity, as indicated by several authors (Walker and Mize, 1982; Mulerji and Kapoor , 1986). Indeed, the highest spore number (132 spores / 100 g soil) was observed in the site Zouala3 where the mycorrhizal intensity of date palm is 2.06%. At the site of Zagoura 2, the number of spores is 90 spores / g soil 100 while the mycorrhizal intensity is 5.7%. According to Jasper et al. (1991), the weak relationship observed between the formation of

endomycorrhizae and quantity of potential propagules encountered can be explained by the fact that the spores are not always viable and sometimes are dormant. In all cases, it is risky to approach the infectious activity of the AMF in a given soil to the number of spores presents in this soil (Diagne and Ingleby, 2003).

Stutz and Morton (1996) emphasized that the relationship between sporulation and colonization of VAM fungi depended on mycorrhizal species, host plants and nutrients in the soil.

Morphological diversity of AM fungi in the studied habitats is supposed to be underestimated and the actual number of endomycorrhizal species could be higher. This underestimation could be due to the small number of soil samples analyzed. Bouamri (2010) reported the presence of 10 species in the date palm rhizosphere of Tafilalet (five species belong to the genus *Glomus*, three *Acaulospora* and two *Scutellospora*). In soils oases of Saudi Arabia, 25 species were detected: 18 species belong to the genus *Glomus*, two species of the genera *Scutellospora* and *Racocetra* and one species of *Acaulospora*, *Paraglomus* and *Ambiospora* (Mohamed Al-Yahya'ei and al., 2011). In the Arabian Peninsula (Arabian desert), Symanczik et al. (2014) presented the characteristics of four species of AMF recovered in the rhizosphere of date palm, namely *Claroideoglomus drummondii*, *Diversispora aurantia* *Diversispora spurca* and *Funneliformis africanum*. In semi-arid areas of Jaipur (India), four genera represented by 11 species have been reported: *Gigaspora*, *Glomus*, *Scutellospora*, *Entrophosphora* and *Sclerocystis* (Sharma et Gheek Batra, 2014). Species of the genus *Gigaspora* are considered best suited for this kind of habitats subject to drought and soil salinity (Muthukumar and Udaiyan, 2002).

This study has highlighted the AMF at the rhizosphere of date palm oasis in the region of Tafilalet and Zagora. The soil of the desert regions rich with endomycorrhizae provides favorable conditions for the growth and development of date palm trees by facilitating their access to minerals and water, and increased tolerance to abiotic stress conditions (drought, salinity of water or soil) and

biotic (attacks of pathogenic microorganisms). The soil of the date palm is probably a reserve of mycorrhizal fungi, may be isolated and used in the restoration of oasis ecosystems and even in improving the production of date palm.

RÉFÉRENCES

- [1] Abohatem M., Zouine J. and El Hadrami I., 2011. Low concentrations of BAP and high rate of subcultures improve the establishment and multiplication of somatic embryos in date palm suspension cultures by limiting oxidative browning associated with high levels of total phenols and peroxidase activities. *Scientia Horticulturae*, 130: 344-348.
- [2] Al-Whaibi M.H. and Khaliel A.S., 1994. The effect of Mg and CA, K and P content of date palm seedlings under mycorrhizal conditions. *Mycoscience*, 35: 213-217.
- [3] Amijie F., Tinker P.B. and Stribley D.P., 1989. The development of endomycorrhizal root systems VII. A detailed study of effects of soil phosphorus on colonizaton.. *New Phytologist*, 111: 435-446
- [4] Anderson R.C., Liberta A.E. and Dickman L.A., 1984. Interaction of vascular plants and vesicular-arbuscular mycorrhizal fungi across a soil moisture-nutrient gradient. *Oecologia*, 64: 111-117.
- [5] Bartschi, H., Gianinazzi-Pearson, V., and I. Vegh 1981. Vesicular-arbuscular mycorrhiza formation and root rot disease (*Phytophthora cinnamomi*) development in *Chamaecyparis lawsoniana*. *Phytopathol. Z.* 102: 213-218.
- [6] Ben Hamid F., 2011. :La filière des dattes communes dans les oasis de Gabés dans le contexte des aléas climatiques et économiques: fonctionnement, atouts et contraintes. Institut national agronomique de Tunisie-Master. <http://www.memoireonline.com/02/12/5304/m-Berta-G-Sampo-S-Gamalero-E-Massa-N-Lemanceau-P.-2005-Suppression-of-Rhizoctonia-root-rot-of-tomato-by-Glomus-mosseae-BEG12-and-Pseudomonas-fluorescens-A6RI-is-associated-with-their-effect-on-the-pathogen-growth-and-on-the-root-morphogenesis.-Eur-J-Plant-Pathol.-111-279-88>.
- [7] Berta G, Sampo S, Gamalero E, Massa N, Lemanceau P., 2005. Suppression of *Rhizoctonia* root-rot of tomato by *Glomus mosseae* BEG12 and *Pseudomonas fluorescens* A6RI is associated with their effect on the pathogen growth and on the root morphogenesis. *Eur J.Plant Pathol.*, 111: 279-88.
- [8] Berta G, Sampo S, Gamalero E, Massa N and Lemanceau P., 2005. Suppression of *Rhizoctonia* root-rot of tomato by *Glomus mossae* BEG12 and *Pseudomonas fluorescens* A6RI is associated with their effect on the pathogen growth and on the root morphogenesis. *Eur. J. Plant Pathol.*, 111: 279-288.
- [9] Binet M.N., Lemoine M.C., Martin C., Chambon C. and Gianinazzi S., 2007. Micropropagation of olive (*Olea europaea* L.) and application of mycorrhiza to improve plantlet establishment. *In Vitro Cellular & Developmental Biology Plant*, 43: 473-478
- [10] Bouammar B., 2010. Le développement agricole dans les régions sahariennes: Etude de cas de la région de Ouargla et de la région de Biskra (2006-2008). Thèse de Doctorat en Sciences économiques. Université KASDI Merbah, Ouargla, Algérie, 296p.
- [11] Bouamri R., Dalpe Y., Serrhini M.N. and Bennani A., 2006. Arbuscular mycorrhizal fungi species associated with rhizosphere of *Phoenix dactylifera* L. in Morocco. *African Journal of Biotechnology*, 5: 6: 510-516.
- [12] Brundrett M., Bougher N., Dell B., Grove T. and Malajczuk N., 1996. Working with Mycorrhizas in Forestry and Agriculture, ACIAR Monograph 32, Australian Centre for International Agricultural Research, Canberra.
- [13] Brundrett M.C., 2009. Mycorrhizal associations and other means of nutrition of vascular plants: understanding the global diversity of host plants by resolving conflicting information and developing reliable means of diagnosis. *Plant Soil*, 320: 37-77.
- [14] Brundrett, M., Melville, L. and Peterson, L. (Editors). 1994. Practical Methods in Mycorrhiza Research. Mycologue Publications, Waterloo, Ont., Canada.
- [15] Diagne O, Ingleby K. 2003. Ecologie des champignons mycorrhiziens arbusculaires infectant *Acacia raddiana*. In *Un Arbre au Désert*, Grouzis M, Floc'h L (eds). IRD Editions: Paris; 205-228.
- [16] Diagne O. and Ingleby K., 2003. Ecologie des champignons mycorrhiziens arbusculaires infectant *Acacia raddiana*. In : Grouzis M. (ed.), *Le Floc'h E. (ed.) : Un arbre au désert : Acacia raddiana*. Paris: IRD, p. 205-228. http://horizon.documentation.ird.fr/exl-doc/pleins_textes/divers10-07/010033301.pdf
- [17] Djerbi M, 1992. Précis de phoeniciculture, F.A.O, Rome, 191 p.
- [18] Djerbi, M., 1982. Le Bayoud en Algérie, Problème et Solution. F.A.O. Regional Projet for palm and Dates Research centre in the Near East and North. Africa, Baghdad Iraq, 45 p.
- [19] Gemma J.N., Koske R.E. and Carreiro M., 1989. Seasonal dynamics of selected species of VA mycorrhizal fungi in a sand dune. *Mycol. Res.*, 92: 317-321.
- [20] Haddouch M., 1993. Situation actuelle et perspectives de développement du palmier dattier au Maroc. In : Ferry M. (ed.), Greiner D. (ed.). *Le palmier dattier dans l'agriculture d'oasis des pays méditerranéens*. Zaragoza : CIHEAM, p. 63-79 (Options Méditerranéennes : Série A. Séminaires Méditerranéens; n. 28).
- [21] Harrier L.A and Watson C.A., 2004. The potential role of arbuscular mycorrhizal (AM) fungi in the bioprotection of plants against soil borne pathogens in organic and/or other sustainable farming systems. *Pest Manag. Sci.*, 60 (2): 149-57.
- [22] Harrison M.J., 1999a. Interfaces and nutrient transport in plant / fungal symbioses. *Journal of Experimental Botany, Special Issues* 50: 1013-102.
- [23] Harrison M.J., 1999b. Biotrophic interfaces and nutrient transport in plant/fungal symbioses *J. Exp. Bot.* (1999) 50 (Special Issue): 1013-1022.
- [24] Husband R., Herre E.A., Turner S.L., Gallery R. and Young J.P., 2002. Molecular diversity of arbuscular mycorrhizal fungi and patterns of host association over time and space in a tropical forest. *Mol. Eco.*: 11(12): 2669-78.

- [25] Jaiti F., Kassami M., Meddich A. and El Hadrami I., 2008. Effect of Arbuscular Mycorrhization on the Accumulation of hydroxycinnamic acid derivatives in date palm seedlings challenged with *Fusarium oxysporum* f. sp. *albedinis*. *Journal of Phytopathology*, 156 (11-12): 641–646.
- [26] Jasper D.A., Abbot L.K. and Robson A.D., 1991. The effect of soil disturbance on
- [27] Jasper, D.A, Robson A.D. and Abbott L.K., 1987. The effect of surface mining on the infectivity of vesicular arbuscular mycorrhizal fungi. *Australian Journal of Botany* 35: 642-652.
- [28] Kaushal S., 2000. Influence of edaphic factors on VAMF spore population and root colonization in *Acacia nilotica* in Rajasthan. *J.Mycol. Plant Pathol.*, 30: 386-388.
- [29] Khaliel A.S. and Abou-Heilah A.N., 1985. Formation of vesicular-arbuscular mycorrhizae in *Phoenix dactylifera* L., cultivated in Qassim region, Saudi Arabia. *Pak. J. Bot.* 17 (2): 267-270.
- [30] Killian, C. et Maire, R., 1930. Le Bayoud, maladie du dattier. *Bull. Soc. Hist. Nat. Agr.*, 21: 89-101.
- [31] Koske R.E., 1987. Distribution of VA mycorrhizal fungi along a latitudinal temperature gradient. *Mycologia*, 79: 55-68
- [32] Labidi S., Ben Jeddi F., Tisserant B., Debiante D., Rezgui S., Grandmougin-Ferjani A. and Lounès-Hadj Sahraoui, 2012. Role of arbuscular mycorrhizal symbiosis in root mineral uptake under CaCO₃ stress. *Mycorrhiza*, 22: 337-345
- [33] Labidi S., Calonne M., Jeddi F.B., Debiante D., Rezgui S. Laruelle F., Tisserant B., Grandmougin-Ferjani A. and Sahraoui, A.L.H., 2011. Calcareous impact on arbuscular mycorrhizal fungus development and on lipid peroxidation in monoxenic roots. *Phytochemistry*, 72 (18): 2335-2341.
- [34] Labidi, S., Nasr H., M. Zouaghi M. and H. Wallander H., 2007. Effects of compost addition on extra-radical growth of arbuscular mycorrhizal fungi in *Acacia tortilis* ssp. *raddiana* savanna in a pre-Saharan area. *Appl. Soil Ecol.*, 35:184-192.
- [35] Lagrange A., 2009. Études écologiques et microbiologiques des espèces du genre *Costularia* (Cyperaceae), pionnières des sols ultramafiques en Nouvelle-Calédonie : perspectives d'application à la restauration écologique. Thèse de Doctorat, Université de la Nouvelle-Calédonie, 234p.
- [36] Mahesh V., 2002. Studies on arbuscular mycorrhizal fungi associated with some grasses in industrially polluted soil and its influence on effluent of tolerance of *Sorghum bicolor* (L.) Moench., P D Thesi, Bharathidasan, University, Tiruchirapali, Tamil Nadu, India, 155 p
- [37] Marschner H. and Cakmak I., 1986. Mechanism of phosphorus-induced zinc deficiency in cotton. *Physiologia Plantarum*, 68: 491-96.
- [38] Mohamed N. Al-Yahya'ei, Fritz Oehl, Vallino M., Lumini E. Redecker D., Wiemken A. and Bonfante P., 2011. Unique arbuscular mycorrhizal fungal communities uncovered in date palm plantations and surrounding desert habitats of Southern Arabia. *Mycorrhiza*, 21,(3): 195-209
- [39] Mohammad M.J., Hamad S.R. and Malkawi H.I., 2003. Population of arbuscular mycorrhizal fungi in semi-arid environment of Jordan as influenced by biotic and abiotic factors. *J. Arid Environ.* 53: 409-417.
- [40] Morte A. and Honrubia M., 2002. Growth response of *Phoenix canariensis* to inoculation with arbuscular mycorrhizal fungi. *Palms*, 46: 76–80.
- [41] Muthukumar T. and Udaiyan K., 1999. Spore-in-spore syndrome in vesicular mycorrhizal fungi and its seasonality in a tropical grassland. *Nova Hedwigia*, 68: 339–49.
- [42] Muthukumar T. and Udaiyan K., 2000. Arbuscular mycorrhizas of plants growing in the Western Ghats region, Southern India. *Mycorrhiza*, 9:297–313
- [43] Muthukumar T. and Udaiyan K., 2002. Arbuscular mycorrhizal fungal composition in semi-arid soils of Western Ghats, southern India: *Curr. Sci*, 82 (6): 25: 624-628.
- [44] Muthukumar T., Sha L., Yang X., Cao M., Tang J. and Zheng Z., 2003. Mycorrhiza of plants in different vegetation types in tropical ecosystems of Xishuangbanna, southwest China. *Mycorrhiza* 13: 289–97.
- [45] Muthukumar T., Udaiyan K. and Manian S., 1994. Vesicular-arbuscular mycorrhizal fungi in Western Ghats. *Indian Botanical Contact*, 10 (2): 79-83.
- [46] Nehila A., Bekki A. and Ighil-Hariz Z., 2012. Rôle de la symbiose mycorrhizienne dans la tolérance aux stress abiotiques. Second edition of the international congress: "Microbial Biotechnology for Development"(MICROBIOD 2), 02-04 October 2012, web site: www.uca.ma/microbiona
- [47] Oihabi A., 1991. Étude des endomycorhizes à vésicules et arbuscules sur le Bayoud et la nutrition du palmier dattier. Thèse de doctorat d'État, Marrakech, Maroc, 117 p.
- [48] Phillips J. M. and D. S. Hayman D., 1970. Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Transactions of the British Mycological Society*, 55: 157-160.
- [49] Radi M., Meddich A., Duponnois R. et Hafidi M., 2009. Rôle des champignons mycorrhiziens à vésicules et à arbuscules MVA des sols de la Ville de Marrakech dans l'amélioration de la croissance et du développement du palmier dattier (*Phoenix dactylifera* L.) soumis à un déficit hydrique. Congrès international: "Biotechnologie Microbienne au service du Développement" (MICROBIOD – 2009) 02-05 Novembre 2009. Université Cadi Ayyad, Marrakech, Maroc, www.ucam.ac.ma/microbiona.
- [50] Radi M.; Meddich A.; Duponnois R.; Hafidi M. Université Cadi Ayyad, Marrakech, Maroc Congrès international: "Biotechnologie Microbienne au service du Développement" (MICROBIOD – 2009) 02-05 Novembre 2009, site web: www.ucam.ac.ma/microbiona

- [51] Saaidi M., 1992. Comportement au champ de 32 cultivars de palmier dattier vis-à-vis du bayoud: 25 années d'observations. *Agronomie*, 12: 359-70.
- [52] Sbiai A., 2011. Matériaux composites à matrice époxyde chargée par des fibres de palmier dattier : effet de l'oxydation au tempo sur les fibres. Thèse de Doctorat, Institut National des Sciences Appliquées de Lyon, 250p
- [53] Schenck N.C. and Smith G.S., 1982. Additional new and unreported species of mycorrhizal fungi (Endogonaceae) from Florida. *Mycologia*, 74: 77-92.
- [54] Schüßler A, Schwarzott D, Walker C, 2001. A new fungal phylum, the Glomeromycota : phylogenese and evolution. *Mycol Res.*, 105 :1413-21.
- [55] Sedra, My.H., 1985. Potentiel infectieux et réceptivité de quelques sols de palmeraie à la fusariose vasculaire du palmier dattier (Bayoud) causé par *Fusarium oxysporum* f.sp. *albedinis*. Thèse de 3ème cycle agronomie. IAV. Hassan II. Rabat, Maroc, 99p.
- [56] Senoussi A., 1999. Gestion de l'Espace Saharien en Algérie: Symbiose ou Confrontation entre systèmes productifs en milieu agricole et pastoral? « Cas de la Région de Ouargla». Thèse de Doctorat de l'Université Toulouse II, France, 404p.
- [57] Sharma N. and Neha Gheek Batra N.G., 2014. Diversity of arbuscular mycorrhizal fungi associated with the rhizosphere of *Phoenix dactylifera* L. in semi-arid soils. *Int. J. Pharm. Bio. Sci.*, 5(1): 819 – 826
- [58] Sieverding E., 1991. Vesicular-arbuscular Mycorrhiza Management in Tropical Agrosystems. GTZ, Eschborn, Germany, 371p.
- [59] Souna F., A. Chafi, Chakroune K., Himri I., Bouakka M. and Hakkou A., 2010. Effect of mycorrhization and compost on the growth and the protection of date palm (*Phoenix dactylifera* L.) against Bayoud disease. *American-Eurasian Journal of Sustainable Agriculture*, 4(2) : 260-267
- [60] Strullu D.G. and Plenchette C., 1991. Les mycorrhizes en horticulture. *PHM Revue Horticole*, 352: 50-55
- [61] Strullu D.G., 1990. Les mycorrhizes des arbres et plantes cultivées. Collection TEC & DOC, Lavoisier, Paris, 250 p.
- [62] Stutz J.C. and Morton J.B., 1996. Successive pot cultures reveal high species richness of arbuscular endomycorrhizal fungi in arid ecosystem. *Canadian Journal of Botany*, 74: 1883-1889.
- [63] Symanczik S., Błaskowski J., Sally Koegel S., Boller T., Wiemken A. and Al-Yahya'ei M. N., 2014. Isolation and identification of desert habituated arbuscular mycorrhizal fungi newly reported from the Arabian Peninsula. *J. Arid. Land*, 6(4): 488-497
- [64] Tinkrr, P. B. 1975. Soil chemistry of phosphorus and mycorrhizal effects on plant growth. In *Endomycorrhizas* (ed. P. E. Sanders, B. Mosse & P. B. Tinker), pp. 353-371, London: Academic Press.
- [65] Trouvelot A., Kough J.L. et Gianinazzi-Pearson V., 1986. Mesure du taux de mycorrhization ayant une signification fonctionnelle. *Dan's: Aspects physiologiques et génétiques des mycorrhizes*, Dijon, 1985. INRA (éd.), pp. 21 7-221.
- [66] Van der Heijden M.A.G., Klironomos J.N. and Ursic M., 1998. Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature*, 396: 69-72.
- [67] vesicular-arbuscular mycorrhizal fungi in soils from different vegetation types. *New Phytologist*, 118: 471-476.
- [68] Walker C. and Sarders F .E., 1986 . Taxonomie concepts in the Endogonaceae :III. The separation of *Sculellospora* gen. nov. from *Giogaspora* Gerdet Trappe. *Mycotaxon*, 27: 169-18 .
- [69] Walker C., 1982. Systematics and taxonomy of arbuscular endomycorrhizal fungi (Glomales) a possible way forward. *Agronomie*, 12: 887-897.
- [70] Wubet T., Weiß M., Kottke I., Teketay D. and Oberwinkler K., 2003. Molecular diversity of arbuscular mycorrhizal fungi in *Prunus africana*, an endangered medicinal tree species in dry Afromontane forests of Ethiopia. *New Phytol.*, 161: 517-52.
- [71] Zeddouk M., 2005. La problématique du développement agricole dans le milieu oasien : cas du Tafilalet. Actes du Symposium International sur le Développement Durable des Systèmes Oasiens du 08 au 10 mars 2005 Erfoud, Maroc - B. Boulanouar & C. Kradi (Eds.). ORMVA du Tafilalet, Errachidia, Maroc, pp. 635-645.
- [72] Zegaye F., Khalid A., Hasnaoui A., Caid Serghini H. and El Amrani A., 2012. Ectomycorrhization of Date Palm and Carob Plants *Global Journal of Science Frontier Research Biological Sciences*, 12 (8): 1-9.
- [73] Zhao Z.W., Xia Y.M., Qin X.Z., Li X.W., Cheng L.Z., Sha T. and Wang G.H., 2001. Arbuscular mycorrhizal status of plants and the spore density of arbuscular mycorrhizal fungi in the tropical rainforest of Xishuangbanna, South China. *Mycorrhiza*, 11: 159-62.
- [74] Zougari-Elwedi B., Sanaa M., Labidi S. et Lounès-Haj Sahraoui A., 2012. Evaluation de l'impact de la mycorrhization arbusculaire sur la nutrition minérale des plantules de palmier dattier. *Etude et Gestion des Sols*, 19(3): 193-202.
